#### **Product Datasheet**

# HER2 Antibody HRP Conjugated

Catalog No: #C00099H

Package Size: #C00099H 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

## Description

Product Name	HER2 Antibody HRP Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	IHC-P IHC-F ICC
Species Reactivity	Hu
Immunogen Description	KLH conjugated synthetic peptide aa 1210-1255 1255 derived from human HER2
Conjugates	HRP
Target Name	HER2
Other Names	NEU; NGL; HER2; TKR1; CD34; HER-2; MLN 19; HER-2 neu; Receptor tyrosine-protein kinase erbB-2;
	Metastatic lymph node gene 19 protein; Proto-oncogene Neu; Proto-oncogene c-ErbB-2; Tyrosine kinase-type
	cell surface receptor HER2; p185erbB2; ERBB2; MLN19
Accession No.	Swiss-Prot#P04626NCBI Gene ID2064
Cell Localization	Cytoplasm
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

## **Application Details**

IHC-P=1:50-200 IHC-F=1:50-200 ICC=1:50-200

#### Background

Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP3 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization. In the nucleus is involved in transcriptional regulation. Associates with the 5'-TCAAATTC-3' sequence in the PTGS2 COX-2 promoter and activates its transcription. Implicated in transcriptional activation of CDKN1A; the function involves STAT3 and SRC. Involved in the transcription of rRNA genes by RNA Pol I and enhances protein synthesis and cell growth.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.