

NKG2D Antibody FITC Conjugated

Catalog No: #C00507F

Package Size: #C00507F 100ul

Description

Product Name	NKG2D Antibody FITC Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	Flow-Cyt IF
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic peptide aa 130-170 215 derived from rat NKG2D
Conjugates	FITC
Target Name	NKG2D
Other Names	NKG2D; Nkrp2; NKG2-D type II integral membrane protein; Killer cell lectin-like receptor subfamily K member 1; NK cell receptor D; NK lectin-like receptor; NKLLR; NKG2-D-activating NK receptor; NKR-P2; CD314; Klrk1
Accession No.	Swiss-Prot#O70215NCBI Gene ID24934
Cell Localization	Extracellular
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Application Details

Flow-Cyt=1:50-200 IF=1:50-200

Background

Function as an activating and costimulatory receptor involved in immunosurveillance upon binding to various cellular stress-inducible ligands displayed at the surface of autologous tumor cells and virus-infected cells. Provides both stimulatory and costimulatory innate immune responses on activated killer (NK) cells, leading to cytotoxic activity. Acts as a costimulatory receptor for T-cell receptor (TCR) in CD8(+) T-cell-mediated adaptive immune responses by amplifying T-cell activation. Stimulates perforin-mediated elimination of ligand-expressing tumor cells. Signaling involves calcium influx, culminating in the expression of TNF-alpha. Participates in NK cell-mediated bone marrow graft rejection. May play a regulatory role in differentiation and survival of NK cells. Binds to ligands belonging to various subfamilies of MHC class I-related glycoproteins.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.