Product Datasheet

Transferrin receptor Antibody FITC Conjugated

Catalog No: #C00528F

Package Size: #C00528F 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

| Product Name | Transferrin receptor Antibody FITC Conjugated |
|-----------------------|---|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Purification | Purified by Protein A. |
| Applications | Flow-Cyt |
| Species Reactivity | Hu Ms Rt |
| Immunogen Description | KLH conjugated synthetic peptide aa 620-670 760 derived from human Transferrin receptor |
| Conjugates | FITC |
| Target Name | Transferrin receptor |
| Other Names | T9; TR; TFR; p90; CD71; TFR1; TRFR; Transferrin receptor protein 1; TFRC |
| Accession No. | Swiss-Prot#P02786NCBI Gene ID7037 |
| Cell Localization | Extracellular |
| Concentration | 1mg ml |
| Formulation | 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. |
| Storage | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |

Application Details

Flow-Cyt=2ug/Test

Background

Cellular uptake of iron occurs via receptor-mediated endocytosis of ligand-occupied transferrin receptor into specialized endosomes. Endosomal acidification leads to iron release. The apotransferrin-receptor complex is then recycled to the cell surface with a return to neutral pH and the concomitant loss of affinity of apotransferrin for its receptor. Transferrin receptor is necessary for development of erythrocytes and the nervous system (By similarity). A second ligand, the heditary hemochromatosis protein HFE, competes for binding with transferrin for an overlapping C-terminal binding site.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.