

c-Kit Antibody FITC Conjugated

Catalog No: #C00532F

Package Size: #C00532F 100ul

Description

Product Name	c-Kit Antibody FITC Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	Flow-Cyt ICC IF
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic peptide aa 360-405 976 derived from human c-Kit
Conjugates	FITC
Target Name	c-Kit
Other Names	PBT; SCFR; C-Kit; CD117; Mast stem cell growth factor receptor Kit; Piebald trait protein; Proto-oncogene c-Kit; Tyrosine-protein kinase Kit; p145 c-kit; v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog; KIT
Accession No.	Swiss-Prot#P10721NCBI Gene ID3815
Cell Localization	Extracellular
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Application Details

Flow-Cyt=1:50-200 ICC=1:50-200 IF=1:50-200

Background

Tyrosine-protein kinase that acts as cell-surface receptor for the cytokine KITLG SCF and plays an essential role in the regulation of cell survival and proliferation, hematopoiesis, stem cell maintenance, gametogenesis, mast cell development, migration and function, and in melanogenesis. In response to KITLG SCF binding, KIT can activate several signaling pathways. Phosphorylates PIK3R1, PLCG1, SH2B2 APS and CBL. Activates the AKT1 signaling pathway by phosphorylation of PIK3R1, the regulatory subunit of phosphatidylinositol 3-kinase. Activated KIT also transmits signals via GRB2 and activation of RAS, RAF1 and the MAP kinases MAPK1 ERK2 and or MAPK3 ERK1. Promotes activation of STAT family members STAT1, STAT3, STAT5A and STAT5B. Activation of PLCG1 leads to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate. KIT signaling is modulated by protein phosphatases, and by rapid internalization and degradation of the receptor. Activated KIT promotes phosphorylation of the protein phosphatases PTPN6 SHP-1 and PTPRU, and of the transcription factors STAT1, STAT3, STAT5A and STAT5B. Promotes phosphorylation of PIK3R1, CBL, CRK (isoform Crk-II), LYN, MAPK1 ERK2 and or MAPK3 ERK1, PLCG1, SRC and SHC1.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.