

5-HTR2 Antibody FITC Conjugated

Catalog No: #C00810F

Package Size: #C00810F 100ul

Description

Product Name	5-HTR2 Antibody FITC Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	Flow-Cyt,IF
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic peptide aa 80-130 471 derived from human 5-HTR2A
Conjugates	FITC
Target Name	5-HTR2
Other Names	HTR2; 5-HT2A; 5-hydroxytryptamine receptor 2A; 5-HT-2; 5-HT-2A; Serotonin receptor 2A; HTR2A
Accession No.	Swiss-Prot#P28223NCBI Gene ID3356
Cell Localization	Cytoplasm
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Application Details

Flow-Cyt=2ug/Test IF=1:50-200

Background

G-protein coupled receptor for 5-hydroxytryptamine (serotonin). Also functions as a receptor for various drugs and psychoactive substances, including mescaline, psilocybin, 1-(2,5-dimethoxy-4-iodophenyl)-2-aminopropane (DOI) and lysergic acid diethylamide (LSD). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors. Beta-arrestin family members inhibit signaling via G proteins and mediate activation of alternative signaling pathways. Signaling activates phospholipase C and a phosphatidylinositol-calcium second messenger system that modulates the activity of phosphatidylinositol 3-kinase and promotes the release of Ca(2+) ions from intracellular stores. Affects neural activity, perception, cognition and mood. Plays a role in the regulation of behavior, including responses to anxiogenic situations and psychoactive substances. Plays a role in intestinal smooth muscle contraction, and may play a role in arterial vasoconstriction.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.