#### **Product Datasheet**

# NMUR1 GPR66 Antibody FITC Conjugated

Catalog No: #C01194F

Package Size: #C01194F 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

### Description

· · · · · · · · · · · · · · · · · · ·	
Product Name	NMUR1 GPR66 Antibody FITC Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	IF
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic peptide derived from human NMUR1
Conjugates	FITC
Target Name	NMUR1 GPR66
Other Names	FM-3; G protein-coupled receptor 66; G-protein coupled receptor 66; G-protein coupled receptor FM-3;
	GPR66; Growth hormone secretagogue receptor family Member 3; Neuromedin U Receptor 1; Neuromedin-U
	receptor 1; NMU-R1; NMUR1; NMUR1_HUMAN.
Accession No.	NCBI Gene ID10316
Cell Localization	Extracellular
Concentration	1mg/ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

## **Application Details**

FCM: 1:20-100

Immunofluorescence: 1:50-200

#### Background

Neuromedin U is a neuropeptide with high activity on smooth muscle. It is widely expressed in gastrointestinal systems and central nervous system (CNS). Peripheral activities of neuromedin U include smooth muscle stimulation, ion transport alterations in the gut and the regulation of local blood flow and adrenocortical function. Neuromedin U receptors 1 and 2 (NMUR1 and NMUR2) are multi-pass membrane proteins that belong to the G-protein coupled receptor 1 family of proteins. Both NMUR1 and NMUR2 act as receptors for the neuromedin U neuropeptide. NMUR1 is detected in peripheral organs, particularly in urogenital and gastrointestinal systems, with highest levels in testis. Ita?s expression in CNS is low, but the protein has been detected in cerebellum, hippocampus, dorsal root ganglia and spinal cord. NMUR2 is predominantly detected in central nervous system with highest levels detected in medulla oblongata, spinal cord and thalamus. It may also be detected in testis but has low levels of expression in peripheral tissues.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.