## **Product Datasheet**

## SEMA6C Antibody FITC Conjugated

Catalog No: #C01241F

Package Size: #C01241F 100ul



Support: tech@signalwayantibody.com

Description SEMA6C Antibody FITC Conjugated **Product Name Host Species** Rabbit Clonality Polyclonal Isotype IgG Purified by Protein A. Purification ICC,IF Applications Species Reactivity Hu Ms Rt Immunogen Description KLH conjugated synthetic peptide derived from human SEMA6C Conjugates **Target Name** SEMA6C Other Names KIAA1869; m Sema Y; m Sema Y2; sema domain, transmembrane domain TM, and cytoplasmic domain, semaphorin 6C; Sema Y antibody Semaphorin 6C [Precursor]; Semaphorin Y; SEMAY; SEM6C\_HUMAN.

0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

## **Application Details**

Cell Localization

Concentration

Formulation

Storage

ICC=1:50-200 IF=1:50-200

## Background

Semaphorins are a family of cell surface and secreted proteins involved in neural development that are conserved from insects to humans. Members of this family are approximately 750 amino acids in length (including signal sequences) and are defined by a conserved extracellular a?semaphorina? domain of approximately 500 amino acids containing 14-16 cysteines, blocks of conserved sequences and no obvious repeats. The transmembrane semaphorins are characterized by an additional 80 amino acid transmembrane domain and an 80-110 amino acid cytoplasmic domain. SEMA6C, also known as SEMA Y, is a transmembrane protein expressed in fetal brain and adult skeletal muscle. Three isoforms of this semaphorin exist due to alternative splicing: SEMA6C 1, SEMA6C 2 and SEMA6C 3. The extracellular domain of SEMA6C induces growth cone collapse of dorsal root ganglion and plays a role in generation or stability of entorhino-hippocampal synapses.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.

Extracellular

1mg ml