#### **Product Datasheet**

# EN2 Engrailed 2 Antibody FITC Conjugated

Catalog No: #C01298F

Package Size: #C01298F 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

### Description

| Product Name          | EN2 Engrailed 2 Antibody FITC Conjugated   |
|-----------------------|--|
| Host Species          | Rabbit   |
| Clonality             | Polyclonal   |
| Isotype               | IgG  |
| Purification          | Purified by Protein A.   |
| Applications          | ICC IF   |
| Species Reactivity    | Hu Ms Rt   |
| Immunogen Description | KLH conjugated synthetic peptide aa 300-333 333 derived from human EN2 Engrailed 2 |
| Conjugates            | FITC   |
| Target Name           | EN2 Engrailed 2  |
| Other Names           | Homeobox protein engrailed-2; Homeobox protein en-2; Hu-En-2; EN2                  |
| Accession No.         | Swiss-Prot#P19622NCBI Gene ID2020  |
| Cell Localization     | Nucleus  |
| Concentration         | 1mg ml   |
| Formulation           | 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.                   |
| Storage               | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.    |

### **Application Details**

ICC=1:50-200 IF=1:50-200

## Background

The engrailed-2 gene, EN2, a murine homolog of the Drosophila homeobox gene engrailed (EN), is required for midbrain and cerebellum development and dorsal ventral patterning of the limbs as well as apical ectodermal ridge formation. In Drosophila, the EN gene plays an important role during development in segmentation, where it is required for the formation of posterior compartments. Human EN-1 and EN-2 are homeodomain-containing proteins and have been implicated in the control of pattern formation during development of the central nervous system. Different mutations in the mouse homologs, EN-1 and EN-2, produce different developmental defects that frequently are lethal. EN-1 is highly expressed by essentially all dopaminergic neurons in the substantia nigra and ventral tegmentum. EN-1 and EN-2 regulate expression of a-synuclein, a gene that is genetically linked to Parkinsona?s disease. During early brain development mouse EN-2 is expressed in a broad band across most of the mid-hindbrain region. EN-2 is also expressed in mouse myoblasts and has been assiciated with cerebellar hypoplasia.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.