## **Product Datasheet**

## Frizzled 9 CD349 Antibody PE Conjugated

Catalog No: #C01531P

Package Size: #C01531P 100ul



Support: tech@signalwayantibody.com

Description

Product Name	Frizzled 9 CD349 Antibody PE Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	ICC IF
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic peptide aa 180-229 591 derived from human Frizzled 9 CD349
Conjugates	PE
Target Name	Frizzled 9 CD349
Other Names	FZD3; CD349; Frizzled-9; Fz-9; hFz9; FzE6; FZD9
Accession No.	Swiss-Prot#O00144NCBI Gene ID8326
Cell Localization	Extracellular
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

## **Application Details**

ICC=1:50-200 IF=1:50-200

## Background

Receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes. A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and or in differentiated tissues.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.