

HIC1 Antibody HRP Conjugated

Catalog No: #C03431H

Package Size: #C03431H 100ul

Description

Product Name	HIC1 Antibody HRP Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	WB IHC-P IHC-F ICC
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic peptide derived from human HIC1
Conjugates	HRP
Target Name	HIC1
Other Names	Hic 1; Hic-1; Hic1; HIC1_HUMAN; Hypermethylated in cancer 1; Hypermethylated in cancer 1 protein; ZBTB29; Zinc finger and BTB domain-containing protein 29; ZNF901.
Accession No.	NCBI Gene ID3090
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Application Details

WB=1:500-2000 IHC-P=1:50-200 IHC-F=1:50-200 ICC=1:50-200

Background

Hypermethylated in cancer (HIC-1) was originally identified as a target of p53-induced gene expression. HIC-1 is deleted in the genetic disorder Miller-Dieker syndrome (MDS), and the expression of HIC-1 is also frequently suppressed in leukemia and various cancers due to the hypermethylation of specific DNA regions and the resulting transcriptional silencing. These and other studies indicate that HIC-1 acts as a putative tumor suppressor protein that mediates transcriptional repression. HIC-1 is ubiquitously expressed in adult tissues and its structure is defined by five zinc fingers and an N-terminal broad complex POZ (or BTB) domain. In several BTB POZ containing proteins, including BCL-6 and the promyelocytic leukemia zinc-finger (PLZF) oncoprotein, this domain interacts with the SMRT N-CoR-mSin3A HDAC complex and is directly involved in repressing and silencing gene transcription. When this domain is deleted, as with the oncogenic PLZF-RAR chimera of promyelocytic leukemias, this transcriptional repression is attenuated. Conversely, HIC-1 does not interact with components of the HDAC complex, suggesting that HIC-1-induced transcriptional repression is unassociated with the POZ BTB domain.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.