## **Product Datasheet**

## ADH AVP ARVP Antibody FITC Conjugated

Catalog No: #C03804F

Package Size: #C03804F 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

$\overline{}$		4.0	
	escri	nti	าท
$\boldsymbol{\nu}$	COUL	Pur	ווע

Product Name	ADH AVP ARVP Antibody FITC Conjugated	
Host Species	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Purification	Purified by Protein A.	
Applications	IF	
Species Reactivity	Hu	
Immunogen Description	KLH conjugated synthetic peptide derived from human ADH (CYAEDCPRG-NH2)	
Conjugates	FITC	
Target Name	ADH AVP ARVP	
Other Names	Antidiuretic Hormone; Arginine Vasopressin; ADH; Arginine vasopressin neurophysin II; ARVP; AVP; AVP	
	NPII; AVRP; Vasopressin neurophysin 2 copeptin precursor; Vasopressin neurophysin II copeptin; VP.	
Accession No.	NCBI Gene ID551	
Concentration	1mg ml	
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.	
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	

## **Application Details**

IF=1:50-200

## Background

Vasopressin, also known as arginine vasopressin (AVP) or antidiuretic hormone (ADH), is a posterior pituitary hormone that is synthesised in the hypothalamus. Vasopressin is synthesised as a precursor protein that consists of arginine vasopressin and two associated proteins, neurophysin 2 and the glycopeptide copeptin. Vasopressin, together with its carrier protein neurophysin II, is packaged into neurosecretory vesicles and transported axonally to the nerve endings in the neurohypophysis, where it is either stored or secreted into the bloodstream. Vasopressin acts as a growth factor by enhancing pH regulation through acid-base transport systems. It has a direct antidiuretic action on the kidney and also causes vasoconstriction of the peripheral vessels. Vasopressin can also contract smooth muscle during parturition and lactation. It also plays a role in cognition, tolerance, adaptation and complex sexual and maternal behaviour, as well as in the regulation of water excretion and cardiovascular functions. Mutations in the vasopressin precursor cause autosomal dominant neurohypophyseal diabetes insipidus (ADNDI), which is characterised by persistant thirst, polydipsia and polyuria.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.