#### **Product Datasheet**

# Histone H3.3(succinyl-K79) Antibody

Catalog No: #SAB497S

Package Size: #SAB497S 50ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

### Description

Product Name	Histone H3.3(succinyl-K79) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were
	purified by affinity-chromatography using epitope-specific peptide.
Applications	Custom antibody
Species Reactivity	Hu
Immunogen Type	Peptide-KLH
Conjugates	Unconjugated
Target Name	H3-3A
Other Names	H3.3A, H3F3, H3F3A
Accession No.	uniprot:P84243
Calculated MW	15kDa
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%
	sodium azide and 50% glycerol.
Storage	Store at $-200\Omega\frac{1}{2}$ o $\Omega\frac{1}{2}$ C for long term preservation (recommended). Store at $40\Omega\frac{1}{2}$ o $\Omega\frac{1}{2}$ C for short term use.

## **Application Details**

Western blotting: 1:500~1:1000

## Background

Variant histone H3 which replaces conventional H3 in a wide range of nucleosomes in active genes. Constitutes the predominant form of histone H3 in non-dividing cells and is incorporated into chromatin independently of DNA synthesis. Deposited at sites of nucleosomal displacement throughout transcribed genes, suggesting that it represents an epigenetic imprint of transcriptionally active chromatin. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.